

Quartiles, Deciles and Percentiles

Data Science and A.I. Lecture Series

Bindeshwar Singh Kushwaha

PostNetwork Academy

Reach PostNetwork Academy

Website

PostNetwork Academy | www.postnetwork.co

YouTube Channel

www.youtube.com/@postnetworkacademy

PostNetwork Academy Facebook Page

www.facebook.com/postnetworkacademy

LinkedIn

www.linkedin.com/company/postnetworkacademy

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3^{rd} and 7^{th} deciles.

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3^{rd} and 7^{th} deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3^{rd} decile

we will use the position formula:

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between
the 6th and 7th values,
which are 18 and 21

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3^{rd} and 7^{th} deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3^{rd} decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6^{th} and 7^{th} values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6th and 7th values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3rd Decile (D_3) is 18.90.

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3^{rd} and 7^{th} deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3^{rd} decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6^{th} and 7^{th} values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3^{rd} Decile (D_3) is 18.90.

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3^{rd} and 7^{th} deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3^{rd} decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6^{th} and 7^{th} values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3^{rd} Decile (D_3) is 18.90.

To find the 7^{th} decile we will use the position formula:

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3^{rd} and 7^{th} deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3^{rd} decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6^{th} and 7^{th} values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3^{rd} Decile (D_3) is 18.90.

To find the 7^{th} decile we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6th and 7th values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3rd Decile (D_3) is 18.90.

To find the 7th decile we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6th and 7th values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3rd Decile (D_3) is 18.90.

To find the 7th decile we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_7 = \frac{7 \cdot (20+1)}{10} = 14.70$$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between the 6th and 7th values, which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3rd Decile (D_3) is 18.90.

To find the 7th decile we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_7 = \frac{7 \cdot (20+1)}{10} = 14.70$$

D_7 is located between the 14th and 15th values, which are 42 and 45:

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6th and 7th values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3rd Decile (D_3) is 18.90.

To find the 7th decile we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_7 = \frac{7 \cdot (20+1)}{10} = 14.70$$

D_7 is located between the

14th and 15th values,

which are 42 and 45:

$$D_7 = 42 + (0.70 \times (45 - 42))$$

$$D_7 = 42 + 2.1 = 44.1$$

Numerical Example to Compute Deciles of Ungrouped Data

Given the data set 3, 6, 9, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57 and 60, calculate 3rd and 7th deciles.

Computing Deciles

Given data set: 3, 6, 9, ..., 60

To find the 3rd decile

we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_3 = \frac{3 \cdot (20+1)}{10} = 6.30$$

D_3 is located between

the 6th and 7th values,

which are 18 and 21

$$D_3 = 18 + (0.3 \times (21 - 18))$$

$$D_3 = 18 + 0.9 = 18.90$$

Thus, the 3rd Decile (D_3) is 18.90.

To find the 7th decile we will use the position formula:

$$\text{Position of } D_k = \frac{k \cdot (n+1)}{10}$$

where $n = 20$

$$\text{Position of } D_7 = \frac{7 \cdot (20+1)}{10} = 14.70$$

D_7 is located between the

14th and 15th values,

which are 42 and 45:

$$D_7 = 42 + (0.70 \times (45 - 42))$$

$$D_7 = 42 + 2.1 = 44.1$$

Thus, the 7th Decile D_7 is 44.10

← Values →







