## Deterministic to Random: The Role of Probability in AI and Data Science

Data Science and A.I. Lecture Series

Bindeshwar Singh Kushwaha

PostNetwork Academy

Bindeshwar Singh Kushwaha (PostNetwork Academy) Deterministic to Random: The Role of Probability in Al and Data S

• An **experiment** refers to an operation or activity that can produce some well-defined outcome(s).

E nar

イロト イヨト イヨト イヨト

- An **experiment** refers to an operation or activity that can produce some well-defined outcome(s).
- Types of experiments:

- An **experiment** refers to an operation or activity that can produce some well-defined outcome(s).
- Types of experiments:
  - Deterministic Experiments

- An **experiment** refers to an operation or activity that can produce some well-defined outcome(s).
- Types of experiments:
  - Deterministic Experiments
  - Random (or Probabilistic) Experiments

3

イロン 不得 とくほ とくほ とう

• These are experiments that have a fixed outcome or result, no matter how many times they are repeated under the same conditions.

- These are experiments that have a fixed outcome or result, no matter how many times they are repeated under the same conditions.
- Example: The Sum of Angles in a Triangle

- These are experiments that have a fixed outcome or result, no matter how many times they are repeated under the same conditions.
- Example: The Sum of Angles in a Triangle
  - $\bullet\,$  In a plane, for any triangle chosen, the sum of its angles is always  $180^\circ.$

- These are experiments that have a fixed outcome or result, no matter how many times they are repeated under the same conditions.
- Example: The Sum of Angles in a Triangle
  - $\bullet\,$  In a plane, for any triangle chosen, the sum of its angles is always  $180^\circ.$
  - This is true regardless of the shape or size of the triangle (e.g., equilateral, isosceles, or scalene).

- These are experiments that have a fixed outcome or result, no matter how many times they are repeated under the same conditions.
- Example: The Sum of Angles in a Triangle
  - $\bullet\,$  In a plane, for any triangle chosen, the sum of its angles is always  $180^\circ.$
  - This is true regardless of the shape or size of the triangle (e.g., equilateral, isosceles, or scalene).
  - This result can be derived mathematically and verified experimentally by measuring angles.

- These are experiments that have a fixed outcome or result, no matter how many times they are repeated under the same conditions.
- Example: The Sum of Angles in a Triangle
  - $\bullet\,$  In a plane, for any triangle chosen, the sum of its angles is always  $180^\circ.$
  - This is true regardless of the shape or size of the triangle (e.g., equilateral, isosceles, or scalene).
  - This result can be derived mathematically and verified experimentally by measuring angles.
- In science and engineering, deterministic experiments ensure consistency and reliability of results.

- These are experiments that have a fixed outcome or result, no matter how many times they are repeated under the same conditions.
- Example: The Sum of Angles in a Triangle
  - $\bullet\,$  In a plane, for any triangle chosen, the sum of its angles is always  $180^\circ.$
  - This is true regardless of the shape or size of the triangle (e.g., equilateral, isosceles, or scalene).
  - This result can be derived mathematically and verified experimentally by measuring angles.
- In science and engineering, deterministic experiments ensure consistency and reliability of results.
  - Example: Testing Ohm's Law under the same conditions always results in the same relationship between voltage, current, and resistance.

• These experiments do not produce the same outcome every time, even when repeated under identical conditions.

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.

3

イロト 不同 トイヨト イヨト

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.

3

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.

э.

イロト 不得 トイヨト イヨト

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:

3

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:
    - Rolling a fair six-sided die can produce any number from 1 to 6.

3

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:
    - Rolling a fair six-sided die can produce any number from 1 to 6.
    - $\bullet\,$  Each outcome has an equal probability of 1/6.

3

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:
    - Rolling a fair six-sided die can produce any number from 1 to 6.
    - Each outcome has an equal probability of 1/6.
    - The unpredictability makes this a classic example of a random experiment.

э.

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:
    - Rolling a fair six-sided die can produce any number from 1 to 6.
    - Each outcome has an equal probability of 1/6.
    - The unpredictability makes this a classic example of a random experiment.
  - Drawing a Card from a Shuffled Deck:

э.

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:
    - Rolling a fair six-sided die can produce any number from 1 to 6.
    - Each outcome has an equal probability of 1/6.
    - The unpredictability makes this a classic example of a random experiment.
  - Drawing a Card from a Shuffled Deck:
    - A standard deck contains 52 cards, each of which can be drawn at random.

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:
    - Rolling a fair six-sided die can produce any number from 1 to 6.
    - $\bullet\,$  Each outcome has an equal probability of 1/6.
    - The unpredictability makes this a classic example of a random experiment.
  - Drawing a Card from a Shuffled Deck:
    - A standard deck contains 52 cards, each of which can be drawn at random.
    - $\bullet\,$  The probability of drawing a specific card (e.g., Ace of Spades) is 1/52.

- These experiments do not produce the same outcome every time, even when repeated under identical conditions.
- The outcome in any trial is one of several possible outcomes, which cannot be determined in advance.
- Examples:
  - Tossing a Coin:
    - When tossing a fair coin, the result can be either *heads* or *tails*.
    - $\bullet\,$  The probability of heads is 50%, as is the probability of tails.
    - However, the outcome of each toss cannot be predicted beforehand.
  - Rolling a Die:
    - Rolling a fair six-sided die can produce any number from 1 to 6.
    - $\bullet\,$  Each outcome has an equal probability of 1/6.
    - The unpredictability makes this a classic example of a random experiment.
  - Drawing a Card from a Shuffled Deck:
    - A standard deck contains 52 cards, each of which can be drawn at random.
    - $\bullet\,$  The probability of drawing a specific card (e.g., Ace of Spades) is 1/52.
    - Randomness is ensured by shuffling the deck thoroughly before drawing.

3

イロト 不同 トイヨト イヨト

• Predictive Modeling:

#### • Predictive Modeling:

• Used to estimate the likelihood of future events based on historical data.

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

• Many algorithms, like Naive Bayes, Hidden Markov Models, and probabilistic graphical models, rely heavily on probability theory.

イロト 不得 トイヨト イヨト ニヨー

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

- Many algorithms, like Naive Bayes, Hidden Markov Models, and probabilistic graphical models, rely heavily on probability theory.
- Probabilities help in decision-making under uncertainty.

イロト 不得 トイヨト イヨト ニヨー

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

- Many algorithms, like Naive Bayes, Hidden Markov Models, and probabilistic graphical models, rely heavily on probability theory.
- Probabilities help in decision-making under uncertainty.

#### • Natural Language Processing (NLP):

イロト 不得 トイヨト イヨト ニヨー

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

- Many algorithms, like Naive Bayes, Hidden Markov Models, and probabilistic graphical models, rely heavily on probability theory.
- Probabilities help in decision-making under uncertainty.

#### • Natural Language Processing (NLP):

• Probability is used to model language and predict the likelihood of word sequences.

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

- Many algorithms, like Naive Bayes, Hidden Markov Models, and probabilistic graphical models, rely heavily on probability theory.
- Probabilities help in decision-making under uncertainty.

#### • Natural Language Processing (NLP):

- Probability is used to model language and predict the likelihood of word sequences.
- Example: Predictive text and speech recognition.

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

- Many algorithms, like Naive Bayes, Hidden Markov Models, and probabilistic graphical models, rely heavily on probability theory.
- Probabilities help in decision-making under uncertainty.

#### • Natural Language Processing (NLP):

- Probability is used to model language and predict the likelihood of word sequences.
- Example: Predictive text and speech recognition.

#### • Data Sampling and Hypothesis Testing:

#### • Predictive Modeling:

- Used to estimate the likelihood of future events based on historical data.
- Examples: Fraud detection, customer behavior analysis, and weather forecasting.

#### • Machine Learning Algorithms:

- Many algorithms, like Naive Bayes, Hidden Markov Models, and probabilistic graphical models, rely heavily on probability theory.
- Probabilities help in decision-making under uncertainty.

#### • Natural Language Processing (NLP):

- Probability is used to model language and predict the likelihood of word sequences.
- Example: Predictive text and speech recognition.

#### • Data Sampling and Hypothesis Testing:

• Probability helps in determining confidence intervals and testing the validity of models.

#### Website

PostNetwork Academy | www.postnetwork.co

#### YouTube Channel

www.youtube.com/@postnetworkacademy

#### Facebook Page

www.facebook.com/postnetworkacademy

#### LinkedIn Page

www.linkedin.com/company/postnetworkacademy

イロン 不通 とうせい うけ

# Thank You!